





PubMed

Nucleotide

Protein

Genome

Structure

**PopSet** 

Taxonomy

OMIM

Books

Search

About Entrez

**Text Version** 

**Entrez PubMed** 

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

#### **PubMed Services**

Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

#### Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

**Privacy Policy** 

for

Limits Preview/Index History Clipboard Details

1: TRENDS IN ECOLOGY AND EVOLUTION 2000 Jan;15(1):7-9

Related Articles, Links

ELSEVIER SCIENCE FULL-TEXT ARTICLE

Adaptive cycles: parasites selectively reduce inbreeding in Soay sheep.

O'Brien SJ.

Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD 21702-1201, USA.

PMID: 10603496 [PubMed - as supplied by publisher]

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer

i686-pc-linux-gnu Aug 30 2002 15:17:13

# Open Linking Technologies ARTICLE LOCATOR ELSEVIER SCIENCE DIRECT

# The article you requested:

Adaptive cycles: parasites selectively reduce inbreeding in Soay sheep. S. O'Brien. Trends in Ecology & Evolution, Volume 15, Issue 1, Pages 7-9

has been found in the following Elsevier Science web products. Select a product to retrieve the article from. A subscription is required.



SET ARTICLE LOCATOR OPTIONS

Software and compilation © 2001 Elsevier Science Inc. All rights reserved Feedback



Research Tools Latest

**Reviews** 

Browse

**Journals** 

Browse

Updates Subjects

**Journal** Collection

Journals

News & Comment

My Virtual Search My E-Mail

Reviews

Books & Labware

Science Jobs

Section

Web Links

reviews.bmn.com

Join 🚮 Login 🕮 Feedback 🐠 Help

Join/Login Benefits

Save your searches

# **Trends in Ecology & Evolution**

🔼 Full A-Z Journal List 🕨 Vol. 17, No. 10, October 2002 🕨 Full Text Record

#### Trends in Ecology & Evolution

▶ Latest Issue **Browse Issues** Search this Journal Prices & Subscriptions About this Journal Publishers Site

**Quick Site Search** 

Advanced site search

Adaptive cycles: parasites selectively reduce inbreeding in Soay sheep

Stephen J. O'Brien obrien@ncifcrf.gov Trends in Ecology & Evolution 2000, 15:7-9

Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD 21702-1201, USA

Text only, + thumbnails, + full figures, PDF Publications by Stephen J. O'Brien Related fulltext articles on BioMedNet

**Article Outline** 

Genes and outbreaks Parasites and severe winters A role for genomic diversity Implications and prospects References Copyright

Genes and outbreaks

TRENDS Ecology&

Sometimes, we can predict the consequences of inbreeding or the cost of an infectious disease outbreak on the survival of free-ranging populations, but actually demonstrating the interplay of the two is another thing. The interactions between variant immune defences, parasite load, population demography, and life history parameters (e.g. age, sex and breeding structure) are likely to determine a population's response to new diseases, but tracking and evaluating these parameters has always been an elusive prospect. However, a new study of feral Soay sheep (Ovis aries), in the face of harsh versus mild winters' influence on an endemic fatal nematode, paints a graphic portrait of partitioned fitness components that bear on whether this small island population survives 1. By

simultaneously considering several interacting ecological factors that mediate relative survival or mortality in this population, Coltman et al.  $\frac{1}{2}$  integrate demographic, molecular genetic, and epidemiological analyses to reveal the regulatory influence of individual and population zygosity.


Abrupt reduction in population genetic variation because of inbreeding, population bottlenecks, or genetic drift will usually diminish overall fitness or survival and can lead to extinction 2.3. Infectious diseases are major threats to a species' survival depending on virulence, pathogenesis, mutability and other components that determine the infectious agents' spread, morbidity and mortality. A few examples have demonstrated an advantage for maximum heterozygosity, particularly involving the mammalian major histocompatibility complex (MHC), in the face of exposure to fatal viruses such as HIV-1 (Ref. 4) or hepatitis-B (Ref. 5). Several additional host genes have alleles that can restrict a pathogen's virulence 6. Yet few parameters that monitor the interplay of host population genetic disposition and acute outbreaks have received detailed analysis, mostly because the tools for assessing molecular genetic variation, the knowledge of host restriction genes, and the biological materials collected in field studies during outbreaks have not been widely available. The new Soay sheep study attempts such a task and in the process implicates a critical influence for genome-wide homozygosity - a signal for close inbreeding - on individual sensitivity to parasite dependent mortality, exacerbated during severe winters of food deprivation. By considering the interaction over time of six ecological parameters on population survivorship (Table 1), Coltman et al. have shown that the genomic influence emerges in a fascinating manner.

Modulating co-factor	Sensitive individuals	Resistant individuals
Nongenetic		
Year (climatic extreme)	In 1989, 1992 and 1995	In other years
Age	<1 yr old	>1 yr old
Sex	Males	Females
Survival of previous crash	Naïve	Previous survivors
Density	Those at high density	Those at low density
Antihelminthic therapy	Untreated	Treated
Genetic		
MHC microsatellites	OLADRB-205, -257, -267	OLDRB-263
Genome-wide microsatellites	Those with low heterozygosity	Those with high heterozygosity

#### **Parasites and severe winters**



Soay sheep are a primitive domestic breed introduced to Soay Island in the St Kilda archipelago west of Scotland about 4000 years ago. In 1932, 85 breeding individuals were introduced to the uninhabited Hirta island (638 hectares), where today up to 1800 animals reside. A longitudinal study of population's natural history was initiated in 1985 (by T.H. Clutton Brock, S.D. Albon and P.A. Jewell) and has continued to the present. The Soay sheep are afflicted with virulent strongyle nematode parasites (principally Teladorsagia circumcincta), which damage the gastrointestinal mucosa and result in nutritional deficiencies. The parasite load (determined by fecal egg counts) varies among individuals with little apparent consequence except during severe winters when sheep with high parasite load preferentially succumb to starvation mortality. The population vacillates from boom to bust years in demographic terms (Fig. 1), with harsh winters (1989, 1992 and 1995) causing 40-70% mortality and mild winters promoting rapid population growth. Parasite load in the summer preceding the harsh winters is a strong predictor of mortality for all age classes, although lambs and yearlings are the most susceptible (53% and 36% mortality, respectively). There is also a sex difference in mortality, with males dying at twice the rate of females. The yearly fluctuation in parasite load correlates with population density, whereby adults in more dense population (after good years of reproductive increase) have nearly twice the parasite load, on average, than the same populations following severe winters of high mortality. Treatment of a subset of parasitized sheep (N=121) with antihelminthic therapy abrogated mortality almost completely, linking the parasite's prewinter incidence directly to survival. Each of these nongenetic parameters dramatically influences survivorship, and now the role of the sheep population genetic structure is being factored into the mix.



**Fig. 1.**Annual fluctuations in the Soay sheep (*Ovis aries*) population on Hirta Island (UK; open circles) and in a smaller subset of this study population (Village Bay; filled circles). Severe winters of increasing starvation mortality were evident in 1989, 1992 and 1995 where the sample and entire population showed similar large mortalities (indicated by arrows). *Reproduced, with permission, from <u>Ref. 1</u>.* 

#### A role for genomic diversity



In 1998, Paterson <sup>7</sup> reported that three microsatellite loci within the class II region of the ovine MHC displayed unusually 'even' allele distributions compared with adjacent microsatellite loci outside the MHC. The MHC region microsatellite loci were in strong linkage disequilibrium (the tendency of alleles at linked polymorphic loci to be associated or occur together non-randomly in populations) with each other. The even allele distributions plus the linkage disequilibrium are indicators that powerful selective pressures operate on functional genes within the MHC even today. A subsequent study 8 demonstrated that certain microsatellite alleles within the MHC are associated with parasite load resistance and with juvenile survival, whereas alternative alleles are more prevalent among susceptible sheep with high parasite loads and increased mortality.

These observations are provocative, particularly in light of our growing understanding of the role of natural selection in reinforcing extensive allele heterozygosity at MHC loci as a defence against infecting microbes. MHC class II gene products recognize invading parasites in hosts, present them to T-helper cells, and assist in mounting a humoral immune response against them. That younger lambs (<6 months) in Paterson et al.'s study  $\frac{8}{2}$  failed to show the MHC effects on survival  $\frac{8-10}{2}$  actually supports a role for MHC genes because immune responses depend on exposure and time-dependent immune development.

The new study  $\frac{1}{2}$  on Soay sheep suggests that inbreeding itself, by homogenizing polymorphic loci over the entire genome, produces homozygous lambs that are preferentially eliminated by the parasite-harsh winter combination. Coltman et al. 1 monitored individual genomic heterozygosity by genotyping up to 14 (average 9.7) microsatellite loci and relating diversity to parasite load and survival, but factoring the effects of age, density and winter severity. In nearly every case, genomic homozygosity is associated with higher parasite load and lower survival in harsh winter years. Apparently, the parasite epidemic itself confers selective pressure against homozygosity (inbreeding was actually demonstrated in several individuals), by eliminating less heterozygous sheep preferentially.

When sheep are treated with antihelminthics in the summers preceding harsh winters, overall genome heterozygosity shows no relationship with survival, providing a powerful control population. Untreated animals display genomic-heterozygosity-dependent pathogen resistance, whereas treated animals, apparently relieved of the parasite selective pressure, show no mortality difference among heterozygosity classes. As a consequence of strong selective advantage for increased heterozygosity with every harsh winter, heterozygosity actually increases with age in the surveyed survivors of the population crashes (Fig. 1).

#### **Implications and prospects**





Coltman et al.'s genetic study tracked 1867 sheep over a 10-year interval, providing a valiant attempt to quantify multifactorial genetic and nongenetic regulation of species survival. It is the first study to implicate a fatal parasite outbreak as a selective agent against increased genomic homozygosity, although there have been other examples of recessive genetic influence for specific genes on infectious disease outcomes  $\underline{6}$ . We are used to thinking that genetic drift on islands leads to increased genetic loss and homozygosity; but in these sheep the parasite load actually reverses the process.

There are two remarkable aspects of this study. First Coltman et al.  $\frac{1}{2}$  provide a comprehensive epidemiological analysis of the interaction of genome-wide zygosity with nongenetic parameters that regulate the likelihood of survival in different ways (Fig. 1, Table 1). Second, they demonstrate that the association of survival with genetic heterozygosity vanishes when the parasite selective pressures are eliminated by antihelminthic treatment. It is not often we catch such a glimpse of the consequence of both factor-dependent mortality and acute selective pressures exerted by infectious diseases in an isolated free-ranging natural population.

A puzzle that comes from this study is which of the specific sheep genes are responsible for the heterozygosity relationship? The studies by Paterson et al. 7.8 reveal that the sheep MHC shows signals of historic and recent heterozygote advantage (historic evidence is an excess of nonsynonymous variants among compared MHC class II alleles; recent evidence is the 'even' allele distribution of MHC linked microsatellite loci). Nevertheless, specific MHC loci (e.g. class I, class II, TNF, Tap, or any of a hundred genes in the MHC) were not genotyped and analysed epidemiologically. Somewhat perplexing is the simultaneous association for MHC region and for wholegenome heterozygosity with survival in the same population. Is it possible that the genome-wide homozygosity is a marker for MHC effect? Alternatively, could MHC effects be surrogates for genomewide homozygosity? MHC diversity is frequently used to screen for historic inbreeding in natural

populations. Yet, in at least one case, MHC gene homozygosity influence on infectious disease progression (AIDS) was uncoupled with MHC linked and with overall genome-wide microsatellite zygosity  $\frac{4}{2}$ .

The identification of gene variants that drive these critical population events should soon be possible. With a full-length sequence of both the human and mouse genomes only a few years away, the power of comparative genomics will allow genome scans to pinpoint responsible genes  ${}^{11}$ . A sheep genetic map including integrated coding (Type I) and microsatellite loci is rapidly developing. The Soay sheep population might be one of the first in which interacting population genetic methods, gene-map driven analyses and ecological insight are employed to describe what until recently have been called stochastic variables. Is it possible that these fluctuations can become deterministic as field ecologists harness the power of comparative molecular genetics? It will not be long.

#### References



[1] Coltman D.W. et al. (1999) Parasite-mediated selection against inbred Soay sheep in a free-living, island population.

Evolution, **53**:1259-1267. Cited by

[2] Frankham R. et al. (1995) Conservation Genetics.

Annu. Rev. Genet., 29:305-327. MEDLINE Cited by

[3] Saccheri I. (1998) Inbreeding and extinction in a butterfly metapopulation.

Nature, 392:491-494. Cited by

[4] Carrington M. et al. (1999) HLA and HIV-1: Heterozygote advantage and B\*35-Cw\*04 disadvantage.

Science, 283: 1748-1752. Full text MEDLINE Cited by

[5] Thursz M.R. et al. (1997) Heterozygote advantage for HLA class II type in hepatitis B virus infection.

Nat. Genet., 17:11-12. MEDLINE Cited by

[6] O'Brien S.J. (1998) AIDS: A role for host genes.

Hosp. Pract., 33:53-79. Cited by

[7] Paterson S. (1998) Evidence for balancing selection at the major histocompatibility complex in a free-living ruminant.

J. Hered., 89:289-294. MEDLINE Cited by

[8] Paterson S. et al. (1998) Major histocompatibility complex variation associated with juvenile survival and parasite resistance in a large unmanaged ungulate population (*Ovis aries*, L.)

Proc. Natl. Acad. Sci. U. S. A., 95:3714-3719. Full text MEDLINE Cited by

[9] Stear M.J. et al. (1997) How hosts control worms.

Nature, 389:27. MEDLINE Cited by

[10] Bishop S.C. et al. (1996) Genetic parameters for foecal egg count following mixed, natural, predominantly Ostertagia circumeineta infection and relationships with live weight in young lambs.

Anim. Sci., 63: 423-428.

[11] O'Brien S.J. (1999) The promise of comparative genomics in mammals.

Science, 286:458-481. Full text MEDLINE Cited by

#### Copyright



© 2000Published by Elsevier Science Ltd.

**BioMedNet** 

Research Tools

Reviews

Journal Collection News & Comment Books & Labware Science Jobs Web Links



raised recurrently. Experimental studies of evolution in natural conditions have become feasible in the past ten years, and applications to the evolution of dispersal are eagerly awaited, as is the integrated book<sup>3</sup> financed by the International Union of Biological Sciences (IUBS) and DIVERSITAS that will synthesize results of this Conference.

#### **Acknowledgements**

We thank the meeting organizers and all participants for helping us put this report together.

#### **Regis Ferriere**

Laboratoire d'Ecologie, Ecole Normale Supérieure, 46 rue d'Ulm, 75230 Paris Cedex 05, France (ferriere@biologie.ens.fr)

#### James R. Belthoff

Dept of Biology, Boise State University, Boise, ID 83725, USA (jbeltho@boisestate.edu)

#### Isabelle Olivieri

Institut des Sciences de l'Evolution, Université Montpellier 2, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France (olivieri@isem.isem.univ-montp2.fr)

#### **Sven Krackow**

Verhaltensbiologie, Zoologisches Institut, Universität Zürich-Irchel, 8057 Zürich, Switzerland (skrackow@zool.unizh.ch)

#### References

- 1 Johnson, M.L. and Gaines, M.S. (1990) Evolution of dispersal: theoretical models and empirical tests using birds and mammals. *Annu. Rev. Ecol. Syst.* 21, 449–480
- 2 Dieckmann, U. et al. (1999) The evolutionary ecology of dispersal. *Trends Ecol. Evol.* 14, 88–90
- 3 Clobert, J. et al., eds Dispersal: Individual, Population and Community, Oxford University Press (in press)
- 4 Ronce, O. et al. Perspectives for the study of dispersal evolution. In *Dispersal: Individual*, Population and Community (Clobert, J. et al., eds), Oxford University Press (in press)
- 5 Ferriere, R. et al. How should we define fitness for spatial ecologies? In Dispersal: Individual, Population and Community (Clobert, J. et al., eds), Oxford University Press (in press)

# Adaptive cycles: parasites selectively reduce inbreeding in Soay sheep

C ometimes, we can predict the conse-Dauences of inbreeding or the cost of an infectious disease outbreak on the survival of free-ranging populations, but actually demonstrating the interplay of the two is another thing. The interactions between variant immune defences, parasite load, population demography, and life history parameters (e.g. age, sex and breeding structure) are likely to determine a population's response to new diseases, but tracking and evaluating these parameters has always been an elusive prospect. However, a new study of feral Soay sheep (Ovis aries), in the face of harsh versus mild winters' influence on an endemic fatal nematode, paints a graphic portrait of partitioned fitness components that bear on whether this small island population survives<sup>1</sup>. By simultaneously considering interacting ecological factors that mediate relative survival or mortality in this population, Coltman et al.1 integrate demographic, molecular genetic, and epidemiological analyses to reveal the regulatory influence of individual and population zygosity.

### **Genes and outbreaks**

Abrupt reduction in population genetic variation because of inbreeding, population bottlenecks, or genetic drift will usually diminish overall fitness or survival and can lead to extinction<sup>2,3</sup>. Infectious diseases are major threats to a species' survival depending on virulence, pathogenesis, mutability and other components that determine the infectious

agents' spread, morbidity and mortality. A few examples have demonstrated an advantage for maximum heterozygosity, particularly involving the mammalian major histocompatibility complex (MHC), in the face of exposure to fatal viruses such as HIV-1 (Ref. 4) or hepatitis-B (Ref. 5). Several additional host genes alleles that can restrict pathogen's virulence<sup>6</sup>. Yet few parameters that monitor the interplay of host population genetic disposition and acute outbreaks have received detailed analysis, mostly because the tools for assessing molecular genetic variation, the knowledge of host restriction genes, and the biological materials collected in field studies during outbreaks have not been widely available. The new Soay sheep study attempts such a task and in the process implicates a critical influence for genome-wide homozygosity - a signal for close inbreeding - on individual sensitivity to parasite dependent mortality, exacerbated during severe winters of food deprivation. By considering the interaction over time of six ecological parameters on population survivorship (Table 1). Coltman et al. have shown that the genomic influence emerges in a fascinating manner.

#### Parasites and severe winters

Soay sheep are a primitive domestic breed introduced to Soay Island in the St Kilda archipelago west of Scotland about 4000 years ago. In 1932, 85 breeding individuals were introduced to the uninhabited Hirta island (638 hectares), where

today up to 1800 animals reside. A longitudinal study of population's natural history was initiated in 1985 (by T.H. Clutton Brock, S.D. Albon and P.A. Jewell) and has continued to the present. The Soay sheep are afflicted with virulent strongyle nematode parasites (principally *Telador*sagia circumcincta), which damage the gastrointestinal mucosa and result in nutritional deficiencies. The parasite load (determined by fecal egg counts) varies among individuals with little apparent consequence except during severe winters when sheep with high parasite load preferentially succumb to starvation mortality. The population vacillates from boom to bust years in demographic terms (Fig. 1), with harsh winters (1989, 1992 and 1995) causing 40-70% mortality and mild winters promoting rapid population growth.


Parasite load in the summer preceding the harsh winters is a strong predictor of mortality for all age classes, although lambs and yearlings are the most susceptible (53% and 36% mortality, respectively). There is also a sex difference in mortality, with males dying at twice the rate of females. The yearly fluctuation in parasite load correlates with population density, whereby adults in more dense population (after good years of reproductive increase) have nearly twice the parasite load, on average, than the same populations following severe winters of high mortality. Treatment of a subset of parasitized sheep (N = 121) with antihelminthic therapy abrogated mortality almost completely, linking the parasite's prewinter incidence directly to survival. Each of these nongenetic parameters dramatically influences survivorship, and now the role of the sheep population genetic structure is being factored into the mix.

Modulating co-factor	Sensitive individuals	Resistant individuals
Nongenetic		
Year (climatic extreme)	In 1989, 1992 and 1995	In other years
Age	< 1 yr old	> 1 yr old
Sex	Males	Females
Survival of previous crash	Naive	Previous survivors
Density	Those at high density	Those at low density
Antihelminthic therapy	Untreated	Treated
Genetic		
MHC microsatellites	OLADRB-205, -257, -267	OLDRB-263
Genome-wide microsatellites	Those with low heterozygosity	Those with high heterozygosity

#### A role for genomic diversity

In 1998, Paterson<sup>7</sup> reported that three microsatellite loci within the class II region of the ovine MHC displayed unusually 'even' allele distributions compared with adjacent microsatellite loci outside the MHC. The MHC region microsatellite loci were in strong linkage disequilibrium (the tendency of alleles at linked polymorphic loci to be associated or occur together non-randomly in populations) with each other. The even allele distributions plus the linkage disequilibrium are indicators that powerful selective pressures operate on functional genes within the MHC even today. A subsequent study8 demonstrated that certain microsatellite alleles within the MHC are associated with parasite load resistance and with juvenile survival, whereas alternative alleles are more prevalent among susceptible sheep with high parasite loads and increased mortality.

These observations are provocative, particularly in light of our growing understanding of the role of natural selection in reinforcing extensive allele heterozygosity at MHC loci as a defence against infecting microbes. MHC class II gene products recognize invading parasites in hosts, present them to T-helper cells, and assist in mounting a humoral immune response against them. That younger lambs (<6 months) in Paterson *et al.*'s



**Fig. 1.** Annual fluctuations in the Soay sheep (*Ovis aries*) population on Hirta Island (UK; open circles) and in a smaller subset of this study population (Village Bay; filled circles). Severe winters of increasing starvation mortality were evident in 1989, 1992 and 1995 where the sample and entire population showed similar large mortalities (indicated by arrows). *Reproduced, with permission, from Ref. 1.* 

study<sup>8</sup> failed to show the MHC effects on survival<sup>8–10</sup> actually supports a role for MHC genes because immune responses depend on exposure and time-dependent immune development.

The new study<sup>1</sup> on Soay sheep suggests that inbreeding itself, by homogenizing polymorphic loci over the entire genome, produces homozygous lambs that are preferentially eliminated by the parasite–harsh winter combination.

Coltman *et al.*<sup>1</sup> monitored individual genomic heterozygosity by genotyping up to 14 (average 9.7) microsatellite loci and relating diversity to parasite load and survival, but factoring the effects of age, density and winter severity. In nearly every case, genomic homozygosity is associated with higher parasite load and lower survival in harsh winter years. Apparently, the parasite epidemic itself confers selective pressure against homozygosity (inbreeding was actually demonstrated in several individuals), by eliminating less heterozygous sheep preferentially.

When sheep are treated with antihelminthics in the summers preceding harsh winters, overall genome heterozygosity shows no relationship with survival. providing a powerful control population. Untreated animals display genomicheterozygosity-dependent pathogen resistance, whereas treated animals, apparently relieved of the parasite selective pressure, show no mortality difference among heterozygosity classes. As a consequence of strong selective advantage for increased heterozygosity with every harsh winter, heterozygosity actually increases with age in the surveyed survivors of the population crashes (Fig. 1).

#### Implications and prospects

Coltman *et al.*'s genetic study tracked 1867 sheep over a 10-year interval, providing a valiant attempt to quantify multifactorial genetic and nongenetic regulation of species survival. It is the first study to implicate a fatal parasite outbreak as a selective agent against increased genomic homozygosity, although there have been other examples of recessive genetic influence for specific genes on infectious disease outcomes<sup>6</sup>. We are used to thinking that genetic drift on islands leads to increased genetic loss and homozygosity; but in these sheep the parasite load actually reverses the process.

There are two remarkable aspects of this study. First Coltman *et al.*<sup>1</sup> provide a comprehensive epidemiological analysis of the interaction of genome-wide zygosity with nongenetic parameters that regulate the likelihood of survival in different ways (Fig. 1, Table 1). Second, they demonstrate that the association of survival with genetic heterozygosity vanishes when the parasite

**8** TREE vol. 15, no. 1 January 2000

selective pressures are eliminated by antihelminthic treatment. It is not often we catch such a glimpse of the consequence of both factor-dependent mortality and acute selective pressures exerted by infectious diseases in an isolated free-ranging natural population.

A puzzle that comes from this study is which of the specific sheep genes are responsible for the heterozygosity relationship? The studies by Paterson et al.7,8 reveal that the sheep MHC shows signals of historic and recent heterozygote advantage (historic evidence is an excess of nonsynonymous variants among compared MHC class II alleles; recent evidence is the 'even' allele distribution of MHC linked microsatellite loci). Nevertheless, specific MHC loci (e.g. class I, class II, TNF, Tap, or any of a hundred genes in the MHC) were not genotyped and analysed epidemiologically. Somewhat perplexing is the simultaneous association for MHC region and for whole-genome heterozygosity with survival in the same population. Is it possible that the genome-wide homozygosity is a marker for MHC effect? Alternatively, could MHC effects be surrogates for genome-wide homozygosity? MHC diversity is frequently used to screen for historic inbreeding in natural populations. Yet, in at least one case, MHC gene homozygosity influence on infectious disease progression (AIDS) was uncoupled with MHC linked and with overall genomewide microsatellite zygosity<sup>4</sup>.

The identification of gene variants that drive these critical population events should soon be possible. With a full-length sequence of both the human and mouse genomes only a few years away, the power of comparative genomics will allow genome scans to pinpoint responsible genes<sup>11</sup>. A sheep genetic map including integrated coding (Type I) and microsatellite loci is rapidly developing. The Soay sheep population might be one of the first in which interacting population genetic methods, gene-map driven analyses and ecological insight are employed to describe what until recently have been called stochastic variables. Is it possible that these fluctuations can become deterministic as field ecologists harness the power of comparative molecular genetics? It will not be long.

#### Stephen J. O'Brien

Laboratory of Genomic Diversity, National Cancer Institute, Frederick, MD 21702-1201, USA (obrien@ncifcrf.gov)

#### References

- 1 Coltman, D.W. *et al.* (1999) Parasite-mediated selection against inbred Soay sheep in a free-living, island population. *Evolution* 53, 1259–1267
- 2 Frankham, R. et al. (1995) Conservation Genetics. Annu. Rev. Genet. 29, 305–327
- **3** Saccheri, I. (1998) Inbreeding and extinction in a butterfly metapopulation. *Nature* 392, 491–494
- 4 Carrington, M. et al. (1999) HLA and HIV-1: Heterozygote advantage and B\*35-Cw\*04 disadvantage. Science 283, 1748–1752
- 5 Thursz, M.R. et al. (1997) Heterozygote advantage for HLA class II type in hepatitis B virus infection. Nat. Genet. 17, 11–12
- **6** O'Brien, S.J. (1998) AIDS: A role for host genes. *Hosp. Pract.* 33, 53–79
- 7 Paterson, S. (1998) Evidence for balancing selection at the major histocompatibility complex in a free-living ruminant. *J. Hered.* 89, 289–294
- 8 Paterson, S. et al. (1998) Major histocompatibility complex variation associated with juvenile survival and parasite resistance in a large unmanaged ungulate population (Ovis aries, L.). Proc. Natl. Acad. Sci. U. S. A. 95, 3714–3719
- **9** Stear, M.J. *et al.* (1997) How hosts control worms. *Nature* 389, 27
- 10 Bishop, S.C. et al. (1996) Genetic parameters for foecal egg count following mixed, natural, predominantly Ostertagia circumeineta infection and relationships with live weight in young lambs. Anim. Sci. 63, 423–428
- **11** O'Brien S.J. (1999) The promise of comparative genomics in mammals. *Science* 286, 458–481



# Making the most of your personal subscription

- High quality printouts (from PDF files)
- Links to other articles, other journals and cited software and databases

All you have to do is obtain your subscription key from the address label of your print subscription then:

Go to http://www.trends.com/free\_access.html and click on the large Click Here'button at the bottom of the page.

You will see one of the following:

- (1) A BioMedNet login screen. If you see this, please enter your BioMedNet username and password. If you are not already a member please click on the Toin Now'button and register. Once registered you will go straight to (2) below.
- (2) A box to enter a subscription key. Please enter your subscription key here and click on the Enter 'button.

Once confirmed, go to http://tree.trends.com and view the full-text of TREE

If you get an error page please contact Mike Brown (info@current-trends.com) stating your subscription key and BioMedNet username and password. Please note that you do not need to re-enter your subscription key for TREE, BioMedNet 'remembers' your subscription. Institutional online access is currently only available through Science Direct (http://www.sciencedirect.com) to Science Direct subscribers. There are plans to introduce Institutional access through BioMedNet later this year, further details will be released nearer the time, if you would like to be contacted with this information please contact Customer Services at info@current-trends.com

TREE vol. 15, no. 1 January 2000